

Direct evidence for this is provided by the finding that serum stimulation of quiescent cells induces the phosphorylation of PKC δ (Fig. 3), but not of PKC ζ (8); this parallels the mitogen responsiveness defined by membrane association (19).

PKC isotypes are regulated by allosteric activation. The demonstration here that like cPKCs, both nPKCs and aPKCs are subject to phosphorylation in their activation loop sites, establishes an additional level of physiological control. The definition of the PI 3-kinase/PDK1 pathway leading to this phosphorylation would account for the role of a PI 3-kinase pathway in triggering n/aPKC-dependent responses (9, 10). This pathway may thus operate in concert with the allosteric input to control PKC.

References and Notes

1. J. W. Orr and A. C. Newton, *J. Biol. Chem.* **269**, 27715 (1994).
 2. S. Cazaubon, F. Bornancin, P. J. Parker, *Biochem. J.* **301**, 443 (1994).

3. H. Hug and T. F. Sarre, *ibid.* **291**, 329 (1993).
 4. W. H. Ziegler *et al.*, in preparation.
 5. J. A. Le Good *et al.*, data not shown.
 6. N. T. Goode and P. J. Parker, *FEBS Lett.* **340**, 145 (1994).
 7. A. R. Olivier and P. J. Parker, *Eur. J. Biochem.* **200**, 805 (1991).
 8. J. A. Le Good, W. H. Ziegler, P. J. Parker, unpublished results.
 9. K. Akimoto *et al.*, *EMBO J.* **15**, 788 (1996).
 10. S. Moriya *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **93**, 151 (1996).
 11. A. J. Flint, R. D. Paladini, D. E. Koshland Jr., *Science* **249**, 408 (1990).
 12. P. Garcia-Paramio, Y. Cabrerizo, F. Bornancin, P. J. Parker, *Biochem. J.* **333**, 631 (1998).
 13. D. H. Whelan and P. J. Parker, *Oncogene*, in press.
 14. L. Stephens *et al.*, *Science* **279**, 710 (1998).
 15. L. Stempka *et al.*, *J. Biol. Chem.* **272**, 6805 (1997).
 16. D. R. Alessi *et al.*, *Curr. Biol.* **7**, 776 (1997).
 17. D. R. Alessi, M. T. Kozlowski, Q. P. Weng, N. Morrice, J. Avruch, *ibid.* **8**, 69 (1998).
 18. N. Pullen *et al.*, *Science* **279**, 707 (1998).
 19. A. R. Olivier and P. J. Parker, *J. Biol. Chem.* **269**, 2758 (1994).
 20. Phosphorylation site-specific antisera were generated with oligopeptides seven residues long encompassing the phosphoamino acid at residue 4. The peptides were coupled to keyhole limpet he-

mocyanin and used to immunize rabbits. Sera were characterized for specificity with dephospho- and phospho- forms of the peptide antigen. Protein immunoblotting analysis was carried out in the presence of dephosphopeptide (1 μ g/ml) to suppress immunodetection of the dephosphorylated proteins.

21. S. Stabel, D. Schaap, P. J. Parker, *Methods Enzymol.* **200**, 670 (1991).
 22. D. R. Alessi *et al.*, *Curr. Biol.* **7**, 261 (1997).
 23. T. Maniatis, J. Sambrook, E. F. Fritsch, *Molecular Cloning, a Laboratory Manual* (Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1989).
 24. G. Hansra, F. Bornancin, R. Whelan, B. A. Hemmings, P. J. Parker, *J. Biol. Chem.* **271**, 32785 (1996).
 25. Lysis buffer: 1% (v/v) Triton X-100, 20 mM tris-HCl (pH 7.5), leupeptin (50 μ g/ml), aprotinin (50 μ g/ml), 1 mM dithiothreitol, and 0.1 mM phenylmethylsulfonyl fluoride.
 26. U. K. Laemmli, *Nature* **227**, 680 (1970).
 27. We thank R. Marais (ICR, London) for providing the myc-pEflink vector. We are grateful to the Deutsche Forschungsgemeinschaft (W.H.Z.), to the Medical Research Council (P.C. and D.R.A.), and to The Royal Society (P.C.) for support of this work.

7 April 1998; accepted 17 August 1998

Metapopulation Dynamics, Abundance, and Distribution in a Microecosystem

A. Gonzalez,* J. H. Lawton, F. S. Gilbert, T. M. Blackburn, I. Evans-Freke

The experimental fragmentation of landscapes of a natural ecosystem resulted in declines in the abundance and distribution of most species in the multispecies animal community inhabiting the landscapes and the extinction of many species. These declines caused the deterioration of the positive interspecific relation between local population abundance and distributional extent in this community. However, when patches were connected by habitat corridors, an immigration "rescue effect" arrested declines in both abundance and distribution and maintained the observed positive relation between them. These results demonstrate the importance of metapopulation dynamics and landscape connectivity for the persistence of populations in fragmented landscapes.

The observation that locally abundant species tend to be widespread, whereas locally rare species tend to be narrowly distributed, is one of the most pervasive patterns in ecology (1-3). At present, there are at least eight explanatory hypotheses, ranging from niche breadth (4) to sampling artifact (5), but each lacks conclusive evidence (3, 6). A positive abundance-distribution relation is a prediction of metapopulation theory (1, 3, 7-9). One formulation of this theory, the rescue

effect hypothesis (8), assumes that immigration decreases the probability of a local population becoming extinct (the "rescue effect") and that the rate of immigration per patch increases as the proportion of patches that are occupied increases. This rescue effect results in a positive relation between local abundance and one measure of distribution, namely the number of occupied patches. An important, but hitherto experimentally untested, consequence of this mechanism is that a reduction in the distributional extent of a species, for example, by habitat destruction, will result in a lower per patch immigration rate and thus a decline in the species' local abundance in remaining unaltered patches (10). A further prediction is that reduction or elimination of immigration between habitat patches inhibits the rescue effect and should result

in declines in both abundance and occupancy in patches. We tested the rescue effect hypothesis for an entire animal community using a miniature, moss-based ecosystem. Moss landscapes are ideal because they are easy to manipulate in the field at a scale that is large relative to the size and dispersal abilities of the animal populations that live in them and because they contain an easily sampled, well-known, diverse community of microarthropods living at high densities (11). To date, experimental tests of extinction and immigration processes have been predominantly laboratory-based (12).

The first experiment (13) (Fig. 1A) examined the effects of habitat fragmentation on the distribution and abundance of species in the moss fauna. One year from the start of this experiment, species in nonfragmented control patches exhibited a strong abundance-distribution relation (Fig. 1B). However, in identically sized samples, there was a significant difference in the species richness of control and fragmented communities (14) [means of 17.0 ± 1.03 versus 10.3 ± 0.66 (\pm SEM); $F(1,11) = 27.9$, $P < 0.001$], a decline in mean species richness per patch of 40%. Surviving species still showed a positive relation between abundance and distribution in the fragments (Fig. 1C), but, as predicted by theory (7, 8, 10), the average patch occupancy and abundance of these species had both declined significantly (paired t tests: patch occupancy, $t = 8.43$, $df = 20$, $P < 0.001$; \log_{10} abundance, $t = 4.09$, $df = 20$, $P < 0.001$; Fig. 1D). This experiment demonstrates how large-scale fragmentation of a landscape can result in a near universal decline in both the distribution and the abundance of the species inhabiting that landscape; even species surviving in the fragments declined in abundance,

A. Gonzalez, J. H. Lawton, T. M. Blackburn, National Environment Research Council Centre for Population Biology, Silwood Park, Ascot, Berkshire, SL5 7PY, UK. F. S. Gilbert and I. Evans-Freke, School of Biological Sciences, University Park, Nottingham University, Nottingham NG7 2RD, UK.

*To whom correspondence should be addressed. E-mail: a.gonzalez@ic.ac.uk

REPORTS

exactly as theory predicts (10).

Although the results of the first experiment are exactly as predicted by metapopu-

lation dynamics theory, they may conceivably have been generated by changes in the microclimate of fragmented patches. Accord-

ingly, a second experiment (15) (Fig. 2A) tested the hypothesis that the decline in abundance and patch occupancy observed in the first experiment was due to the reduction or elimination of interpatch movement and consequently the lack of a rescue effect. To test this hypothesis, we again fragmented the landscape but maintained corridors between some habitat patches. We also created "pseudocorridors" to control for the slight increase in area due to the corridors. After 6 months, species in the continuous control patches exhibited a strong abundance-distribution relation. However, 41% of the species present in the isolated fragments and in the pseudocorridor treatments had suffered extinction. Surviving species in the pseudocorridor treatment still showed a significant abundance-distribution relation (Fig. 2B) but declined in both abundance (paired *t* test: $t = 4.98$, $df = 20$, $P < 0.001$; Fig. 2C) and distribution (paired *t* test: $t = 4.50$, $df = 20$, $P < 0.001$) relative to the controls. Species surviving in the pseudocorridor and fragmented treatments did not differ in distribution (paired *t* test: $t = 0.66$, $df = 20$, not significant), but the abundances of species were significantly lower in the pseudocorridor treatment (paired *t* test: $t = -2.46$, $df = 20$, $P = 0.02$). Thus, changes in the moss fauna in the pseudocorridor treatment mirror both those in the completely isolated patch treatment and, importantly, those in the fragmentation treatment of the first experiment. In contrast, only 14.5% of the species in the communities inhabiting the landscapes connected by corridors became extinct. Although extant species in the corridor treatment also showed significant declines in both abundance (paired *t* test: $t = 4.92$, $df = 20$, $P < 0.001$; Fig. 2C) and distribution (paired *t* test: $t = 3.19$, $df = 20$, $P < 0.001$) relative to the controls, these declines were significantly less than those in the pseudocorridor treatment (paired *t* tests: patch occupancy, $t = 2.90$, $df = 20$, $P = 0.009$; \log_{10} abundance, $t = 4.31$, $df = 20$, $P < 0.001$).

These experiments show that, at the landscape scale, habitat fragmentation results in a near universal decline in the distribution and abundance of animal species, with abundances declining even on surviving habitat patches. Exactly as theory predicts, these declines are a consequence of reducing or eliminating immigration between patches in the fragmented landscape. The creation of corridors (but not pseudocorridors of similar area) significantly ameliorates the effects of fragmentation. We suggest that real corridors facilitate dispersal between patches, creating a metacommunity and thereby maintaining the distribution and abundance of species within patches through the rescue effect (16).

A positive abundance-distribution relation is the null expectation if the distribution of




Fig. 1. (A) Schematic representation (not to scale) of a single replicate of the treatment design used in the first experiment (13). (B) The positive relation between abundance and distribution (proportion of patches occupied) for the microarthropods inhabiting the control moss fragments at 12 months [$F(1,28) = 30.3$, $P < 0.001$]. (C) Plot of population density against proportion of patches occupied for the species surviving fragmentation after 12 months (open circles) and for the same species in the control samples (filled circles). For clarity, three control points have been displaced horizontally. (D) Change in population density for the species surviving fragmentation illustrated by plotting population density after fragmentation against density before fragmentation. The 1:1 line has been added to better show that most species decline on fragmented patches. Coincident points are represented on all graphs by larger plot symbols, the largest indicating the coincidence of seven points.




Fig. 2. (A) Schematic representation (not to scale) of the treatment design used in the second experiment (19). (B) The abundance-distribution relation for those species surviving fragmentation in the pseudocorridors (open circles) and for the same species from the control sample communities (filled circles). Coincident points are represented by larger plot symbols. For clarity, several points have been displaced horizontally. (C) The difference in the abundance of each species in the corridor (filled circles) and pseudocorridor (open circles) communities after and before fragmentation. Points below the 1:1 line show a decline due to fragmentation.

REPORTS

individuals follows some stochastic process (5). However, in the absence of immigration, species in fragmented landscapes "slide down" the relation observed in the control communities; many species become extinct and most become rarer. Stochastic processes alone cannot generate these significant differences between treatments. Moreover, the different responses of species to the corridor and pseudocorridor treatments count against hypotheses based on interspecific variation in environmental tolerance. Thus, these results corroborate the rescue effect metapopulation model predicting the near universal relation between abundance and distribution (4, 17). The scale of fragmentation and the dispersal distances of the organisms in this system are likely to be appropriate for observing metapopulation dynamics, and we do not necessarily expect the metapopulation mechanism to pertain at biogeographical scales (4). The results of these experiments emphasize that, at the landscape level, maintaining dispersal is imperative for maintaining the abundance-distribution relation. They also starkly demonstrate how population persistence in the face of habitat fragmentation is critically dependent on the maintenance of landscape connectivity (18).

References and Notes

1. I. Hanski, J. Kouki, A. Halkka, in *Species Diversity in Ecological Communities: Historical and Geographical Perspectives*, R. E. Ricklefs and D. Schluter, Eds. (Univ. of Chicago Press, Chicago, IL, 1993), pp. 108–116.
2. K. J. Gaston, *Oikos* **76**, 211 (1996).
3. ———, T. M. Blackburn, J. H. Lawton, *J. Anim. Ecol.* **66**, 579 (1997).
4. J. H. Brown, *Am. Nat.* **124**, 255 (1984).
5. D. H. Wright, *J. Biogeogr.* **18**, 463 (1991).
6. J. H. Lawton, *Bird Study* **43**, 3 (1996).
7. I. Hanski, *Oikos* **62**, 88 (1991).
8. ———, *Biol. J. Linn. Soc.* **42**, 17 (1991).
9. S. Nee, R. D. Gregory, R. M. May, *Oikos* **62**, 83 (1991).
10. J. H. Lawton, *Trends Ecol. Evol.* **8**, 409 (1993); in *Extinction Rates*, J. H. Lawton and R. M. May, Eds. (Oxford Univ. Press, Oxford, UK, 1995), pp. 147–163.
11. R. C. Davis, *Ecol. Monogr.* **5**, 125 (1981); E. L. Seyd, *Zool. J. Linn. Soc.* **106**, 115 (1992). Experiments were carried out on large, flat, moss-covered limestone rocks near Wirksworth in the Peak District (Derbyshire, UK; OS map reference SK 264566). The moss species present were predominantly *Hypnum cupressiforme*, *Thuidium tamariscinum*, and *Tortella tortuosa*. Within the mosses exists a species-rich microinvertebrate community of producers, herbivores, and predators, including Acari, Collembola, Nematoda, Protista, Rotifera, Tardigrada, algae, bacteria, and fungi. All the organisms in this system have very short generation times, ranging from a few hours to a few months. Here we focus on the microarthropods, mainly Acari and Collembola, which are highly abundant and, in comparison with the other groups, easy to extract and identify. They include herbivores, feeding primarily on fungal hyphae and dead plant material, and predators, feeding on nematodes and other Acari and Collembola. They are susceptible to desiccation and so are highly dependent on the mesic microclimate of their moss habitat.
12. P. H. Warren and K. J. Gaston, *J. Anim. Ecol.* **66**, 730 (1997); M. Holyoak and S. J. Lawler, *ibid.* **65**, 640 (1996); T. V. Burkey, *Am. Nat.* **150**, 568 (1997).
13. Eight experimental moss landscapes, each on a single large rock, were established in a randomized block design. Habitat fragments were created with a template. Moss was scraped away and removed from around the template, leaving fragments of a constant area on bare rock. The template ensured the maintenance of a minimum distance between islands. As in natural isolated fragments, the "inhospitable" habitat between islands is probably a partial rather than an absolute barrier to movement; there is very little evidence concerning the dispersal abilities of microarthropods (19). The species richness and abundance of fragmented and unfragmented communities were compared at regular 2-month intervals over the course of 1 year. This time period is equivalent to many generations for most of the fauna inhabiting this system. Each replicate landscape (Fig. 1A) consisted of six island fragments of moss (20 cm²) (containing populations of 140 to less than 10 individuals) and a larger continuous area of moss acting as a control (at least 50 cm by 50 cm). Every 2 months during the ensuing year, one fragment was randomly chosen and removed from each of the eight replicates. At the same time, an equivalently sized patch was removed from the control area of each replicate. The microarthropod community of all samples was extracted with a Tullgren funnel and sorted into morphotypes. Extraction efficiency was high (14), and the same extraction protocol was applied to all moss samples. About 18,000 individual animals were counted and sorted in total; they were mostly Acari (63%) and Collembola (24%). Species richness was taken as an average of the replicate samples, for both control samples and fragments. The total abundance and distribution of each species across the replicate patches sampled at each time period were also estimated for each treatment. The results here refer to comparisons between control and fragmented communities (30 and 21 species, respectively) made after 12 months in six of the original eight replicates (two others were lost to natural destruction).
14. A. Gonzalez, thesis, University of London (1998).
15. The second experiment was a 6-month repeat of the first, with the addition of two corridor treatments (Fig. 2A). In these two additional treatments, fragmented islands were connected to other fragments by an intact corridor, or a corridor was present but was broken by a small strip (~10 mm) of bare rock. The pseudocorridor treatment accounted for the possible confounding effect of the extra moss area associated with the corridor treatment. The only effect of unbroken corridors, therefore, was to facilitate microarthropod dispersal between moss fragments.
16. F. S. Gilbert, A. Gonzalez, I. Evans-Freke, *Proc. R. Soc. London B* **265**, 1 (1998).
17. K. J. Gaston, *Rarity* (Chapman & Hall, London, 1994).
18. E. O. Wilson and E. Willis, in *Ecology and Evolution of Communities*, M. L. Cody and J. M. Diamond, Eds. (Harvard Univ. Press, Cambridge, MA, 1975), pp. 522–534; P. D. Taylor, K. Fahrig, K. Henein, G. Merriam, *Oikos* **68**, 571 (1993); T. H. Keitt, D. L. Urban, B. T. Milne, *Conserv. Ecol.* [online] **1** (no. 1), 4 (1997). Available at www.consecol.org/journal/vol1/iss1/art4.
19. R. Schenker and W. Block, *Bull. Br. Antarct. Surv.* **71**, 36 (1986).
20. We thank E. Chaneton, M. Bonsall, and J. Birch for helpful comments and discussions on the manuscript; H. Mace for help with a pilot study; J. Murphy for help with mite identification; J. Rieley for identifying the mosses; and M. Wright, O. Petchey, G. Couper, A. Kelley, and B. Morris for help in the field. A.G. was supported by a National Environment Research Council (NERC) studentship. T.M.B. is a Leverhulme Special Research Fellow and was supported in part by NERC grant GST/03/1211.

12 June 1998; accepted 19 August 1998

Mind the gap.

NEW! Science Online's Content Alert Service: With *Science's* Content Alert Service, European subscribers (and those around the world) can eliminate the information gap between when *Science* publishes and when it arrives in the post. This free enhancement to your *Science* Online subscription delivers e-mail summaries of the latest news and research articles published each Friday in *Science* – **instantly**. To sign up for the Content Alert service, go to *Science* Online and eliminate the gap.

Science
www.sciencemag.org

For more information about Content Alerts go to www.sciencemag.org. Click on Subscription button, then click on Content Alert button.